

**Can satellite altimetry and gravimetry data be used to
evaluate Arctic-Ocean satellite sea surface salinity?
– insight from ECCO state estimation**

*Tony Lee, Adam Wang, Severine Fournier, Rom Kwok
NASA Jet Propulsion Laboratory, California Institute of Technology*

Special thanks to Ou Wang of JPL for providing ECCO-v4 model-data differences in the Arctic Ocean and Ichiro Fukumori of JPL for discussion

Motivation, rationales, and approach

Motivation & rationales:

- Changes of Arctic Ocean freshwater are important to ocean, climate, & BGC.
- Sea surface salinity (SSS) data are important to study Arctic freshwater changes, ocean-ice interaction, and interaction with the North Atlantic Ocean.
- Large uncertainties of L-band satellite SSS; but how bad is the S/N ratio?
- Lack of in-situ salinity data to validate satellite SSS.
- SSH from altimetry & OBP from gravimetry can be used to infer column-integrated freshwater changes in the Arctic Ocean (e.g., Morrison et al. 2012)


Questions addressed: To what extent Arctic SSS & SSH-OBP variations are coherent?
Can SSS-OBP data can be used to evaluate satellite SSS?

Significance: validated satellite SSS are complementary to SSH-OBP data to study Arctic freshwater changes due to better sampling (SMOS+Aquarius+SMAP).

Approach: Examine relationships of SSS & SSH-OBP in ECCO-V4 for a proof-of-concept study as a 1st step.


Uncertainty characteristics of L-band Satellite SSS: Tropics & subtropics ✓ high-latitudes ✗

Zonally averaged STD of Δ SSS for (Aquarius V4.0 - Argo-SIO) & (Argo-SIO - Argo-UH)
(Argo-SIO & Argo-UH are two different "ground truth" datasets)



Lee (2016)


SSS signals in the Arctic Ocean are large! So how bad is the S/N ratio for L-band satellite SSS?



Timmermans et al. (2014) and NOAA Arctic Report Card 2013 Update.
Gradient estimates provided by Mike Steele of APL, University of Washington

Comparison of time-mean salinity profile in the Arctic Ocean between Ice-Tethered Profiler data & co-located ECCO estimates


ECCO-v4 reproduces the structure <200 m well, which dominate the steric changes



Courtesy of Ou Wang, JPL


Local correlation of SSS & 0-200 halosteric height: they are coherent over much of the Arctic Ocean

Corr Coef of SSS and Halosteric (200m, With Seasonal Cycle) (1992-2015)



With Seasonal Cycle


Corr Coef of SSS and Halosteric (200m, Without Seasonal Cycle) (1992-2015)



Without Seasonal Cycle


**Local correlation of SSS & SSH:
poor correlation around the Arctic shelves
(because of barotropic coastal Kelvin waves, unrelated to SSS)**

Corr Coef of SSS and SSH (With Seasonal Cycle) (1992-2015)



With Seasonal Cycle


Corr Coef of SSS and SSH (Without Seasonal Cycle) (1992-2015)



Without Seasonal Cycle


**Local correlation of SSS & SSH-OBP:
much strong correlation around the shelves
(because barotropic wave signals have been removed)**

Corr Coef of SSS and SSH-BP (With Seasonal Cycle) (1992-2015)



With Seasonal Cycle

Corr Coef of SSS and SSH-BP (Without Seasonal Cycle) (1992-2015)




Without Seasonal Cycle

**Linear regression slope for SSS vs. SSH-OBP (in psu/cm):
good “calibration” sensitivity around the shelves ~ 0.3 psu/cm**


Regression of SSS and SSH-BP (a0, psu/cm, With Seasonal Cycle) (1992-2015)

Regression of SSS and SSH-BP (a0, psu/cm, Without Seasonal Cycle) (1992-2015)




SSS & SSH-OBP relationships in Chukchi Shelf and Siberian Shelf



Chukchi Shelf



— Normalized SSS anomalies
- - Normalized SSH-OBP anomalies




Siberian Shelf




Standard deviation of Arctic SSS from Aquarius & SMAP: large differences in magnitudes!

**Aquarius
(2011-2015)**



**SMAP
(2015-2017)**



Ongoing work: use altimetry-gravimetry data to evaluate satellite SSS and improve Arctic Ocean SSS retrievals.

Summary

- ECCO ocean-sea ice state estimation product illustrates the strong relationship of SSS and SSH-OBP variations around Arctic shelves.
- This relationship could be used to evaluate satellite SSS in the Arctic Ocean using altimetry-gravimetry data and improve SSS retrievals.